

Validation of Kepler Objects of Interest Stellar Parameters And Eccentricity Distribution from Observed Transit Durations

Peter Plavchan¹, Christopher Bilinski², and Thayne Currie³

ABSTRACT

The announcement of 2300+ candidate transiting exoplanets (KOIs) orbiting ~ 1800 host stars discovered with the Kepler mission enables a plethora of ensemble analysis of the architecture and properties of exoplanetary systems. We use the transit durations to probe the ensemble validity of stellar parameters for Kepler candidate host stars. While the new KOI stellar parameters are an improvement over the previous release, our analysis shows that a systematic trend in the ensemble stellar radii remains, which affects the KOI exoplanet radii inferred from transit depths. We also compare the distribution of observed transit durations to a model generated from the known eccentricities of radial velocity (RV) discovered exoplanets. We find that the Kepler and RV distributions differ at a statistically significant level, even after accounting for the transit probability as a function of the eccentricity, inclination, and periastron angle. Additionally, smaller radii KOIs are preferentially found on more circular orbits. Finally, there is an over-abundance of KOIs with long transit durations. We identify that errors in stellar metallicity and possibly extinction, resulting in under-estimated stellar radii, can account for these sources rather than eccentric planets improbably transiting near apastron.

Subject headings: exoplanets – statistical

1. Introduction

The Kepler mission is revolutionizing our understanding of exoplanets (Borucki et al. 2010), including among its many existing and assuredly forthcoming highlights the discovery of three terrestrial exoplanets orbiting the M dwarf Kepler Object of Interest (KOI) 961 (Muirhead et al. 2012), and a $2.4 R_{\oplus}$ exoplanet in the habitable zone of its host star (Borucki et al. 2012). A list of 312 KOIs was published in Borucki et al. (2011a), derived from Q0-1 Kepler time-series data.

¹NASA Exoplanet Science Institute, California Institute of Technology, M/C 100-22, 770 South Wilson Avenue, Pasadena, CA 91125; plavchan@ipac.caltech.edu

²University of Arizona

³NASA Goddard

This was soon followed by a second list of 1235 KOIs in Borucki et al. (2011b) derived from Q1-Q5 Kepler data. The first two KOI releases relied on stellar parameters from the Kepler Input Catalog (KIC, Brown et al. 2011). Batalha et al. (2012) (hereafter B12) recently announced 2321 candidate transiting exoplanet KOIs orbiting 1783 host stars from an improved pipeline analysis of the Q1-Q6 Kepler data, and provided updated stellar host parameters. This large ensemble of exoplanet candidates, including many multi-exoplanet systems, beckons for the ensemble analysis of exoplanetary system architectures (Fabrycky et al. 2012; Howard et al. 2011; Morton & Johnson 2011; Plavchan & Bilinski 2012; Figueira et al. 2012; Youdin 2011).

The ensemble analysis of KOIs and the determination of the frequency of Earth-sized planets ($\eta_{\oplus}$) in habitable zone orbits relies on the accuracy of the estimated physical stellar host parameters such as mass, radius and effective temperature (Batalha et al. 2010a,b; Pinsonneault et al. 2012; Traub 2012). Recent studies suggest that the Kepler Input Catalog (KIC) mischaracterizes some objects, in particular the stellar surface gravity for stars with effective temperatures less than ~ 4500 K. For example, Mann et al. (2012); Ciardi et al. (2011) demonstrate that M “dwarf” KIC targets brighter than a Kepler magnitude of 14th are most likely giants, and Muirhead et al. (2011) reported the over-estimate of M dwarf radii for KOI host stars from the second KOI candidate release in Borucki et al. (2011b). B12 acknowledges these limitations of the new and old stellar parameters.

In this work we use the transit durations of KOIs for two purposes: one, as a probe of the inferred eccentricity distribution of the candidate exoplanets in comparison to exoplanets discovered with the radial velocity (RV) technique, and two, as a new diagnostic of the accuracy of the estimated stellar host parameters. The analytic framework for these two investigations is already laid out in Ford et al. (2008) (hereafter F08) and Burke (2008) and references therein. A more comprehensive analysis is also already carried out with the Borucki et al. (2011b) KOI list by Moorhead et al. (2011). Additionally, Dawson & Johnson (2012) carry out a thorough analysis when the impact parameter can reliably be estimated for Jovian-sized exoplanet KOIs with high S/N transit time-series. In particular, we confirm that the new KOI stellar parameters and KIC values still have some systematic errors. We also find that the distribution of the KOI transit durations is in disagreement with transit durations modeled from the eccentricity distribution of RV-discovered exoplanets.

2. Samples

We use the second and third tabulation of KOI candidate exoplanet and stellar host parameters from Borucki et al. (2011b) and B12 respectively, the latter of which is currently available at the NASA Exoplanet Archive (<http://exoplanetarchive.caltech.edu/>). Details on the methods used to tabulate these parameters can be found in those references and citations therein. We assume that there are no false-positive KOIs, and for simplicity we treat multiple KOIs orbiting the same host star as independent statistical tests. We make additional subsets of terrestrial, Neptune-like, and

Jovian KOIs at exoplanet radii of $R_{pl} < 2$, $2 < R_{pl} < 6$, and $R_{pl} > 6 R_{\oplus}$ earth radii respectively. We do not filter the KOIs by orbital period or by their host mass, temperature, or radius. We do exclude KOIs from our analysis where only one transit is observed and the tabulated period is only a lower bound.

We also use the eccentricity values of 428 published and confirmed RV-discovered exoplanets as listed at the NASA Exoplanet Archive as of March 7th, 2012, with a mean eccentricity of 0.25. For simplicity, we do not filter the RV exoplanets by planet mass, period, or stellar properties, but we note that both samples are reasonably complete for large planets up to periods of ~ 8 months. Whereas the distribution of periastron angles for RV exoplanets is uniform random on a sphere, the distribution of periastron angles for transiting exoplanets will more strongly favor angles perpendicular to the plane of the sky for eccentric planets due to the increased probability of transit (F08, Burke 2008). Thus, we do not use the measured periastron angles for RV planets since they have an incorrect distribution for comparison to the transiting planets.

3. Observed Transit Duration Anomalies

Equal to τ_0 in F08 in the limit $R_{pl}/R_* \rightarrow 0$, we introduce the transit duration anomaly dimensionless parameter α :

$$\alpha = \frac{aT\pi}{(R_* + R_{pl})P} \quad (1)$$

where α is the ratio of the observed transit duration to the expected transit duration for a circular edge-on orbit, and can be computed from the KOI table parameters in B12. R_* and R_{pl} are the stellar and exoplanet radii respectively, P is the orbital period of the exoplanet, T is the duration of the transit, and a is the semi-major axis of the exoplanet. The semi-major axis of the exoplanet a is in turn calculated from the stellar mass M_* , the orbital period of the exoplanet P , and Kepler’s third law. P and T are observed quantities directly measured from the time-series, whereas R_* and M_* (and consequently a) are estimated from ancillary observations and models such as observed colors, magnitudes, spectroscopy and fits to theoretical isochrones (Batalha et al. 2010a,b, 2012; Muirhead et al. 2011). Values for the transit duration anomaly α are tabulated in Table 1.

For an eccentric, non-edge-on orbit, the square of the transit duration anomaly can be written in a similar fashion to the square of Equation 1 in F08 as:

$$\alpha^2 \approx \left(\frac{d_t}{a}\right)^2 \left(\frac{1}{1-e^2}\right) \left(1 - \frac{b^2}{(1+R_p/R_*)^2}\right) \quad (2)$$

where e is the eccentricity, R_p is the exoplanet radius, b is the impact parameter, and d_t is the exoplanet-star separation during transit as defined in F08. Assuming as in F08 that $a/R_* \gg 1$,

$a/R_p \gg 1$, and $T/P \ll 1$, d_t can be assumed to be approximately constant during transit (hence the approximation rather than equality in Equation 2).

From Equation 2, the eccentricity distribution of transiting exoplanets can be inferred from the distribution of observed transit duration anomalies (e.g. the mean, standard deviation), as discussed in great detail in previous works in Burke (2008); Ford et al. (2008); Moorhead et al. (2011); Barnes (2007). α is the correct quantity to use as a proxy for eccentricity instead of the transit duration excess in hours, since transit durations increase with semi-major axis and period due to slower orbital velocities at a fixed eccentricity.

In particular, we can identify potential eccentric exoplanets with values of $\alpha \ll 1$ transiting near periastron or $\alpha \gg 1$ transiting near apastron. Since b is degenerate with eccentricity, the KOIs with $\alpha \ll 1$ may alternatively be grazing transits. We tabulate these candidates in Table 2. We find in §5 that most KOIs with $\alpha \gg 1$ possess errors in their stellar parameters, and are excluded from Table 2 unless $\log[F_e]/[H] > -0.11$ and $A_V < 0.33$ mag.

The four KOIs with the smallest inferred transit duration anomalies compared to a circular orbit – KOIs 338.01, 338.02, 977.01, and 1054.01 – are reported to orbit giant stars with $\log g < 2.5$ with periods of 1.4–7 days interior to the estimated stellar radii, and may instead be associated with photospheric activity or a blend. Visual inspection of the time-series for KOIs 977 and 1054 support this hypothesis. The KOI 338 system has two candidate exoplanets, exhibits clear transit-like dips that are short w/r/t to the transit period, and is specifically mentioned in B12 and Fabrycky et al. (2012). We also exclude these four KOIs from Table 2.

The mean value for α for all KOIs from B12 is 0.921. The expected value for α assuming all exoplanets are on circular orbits with a random impact parameter between 0 and 1 with $\langle b \rangle = 0.5$ is $\sqrt{3}/2 \sim 0.866$. The higher and statistically significant observed mean value of α suggests that there exists a systematic ensemble under-(over-)estimate of stellar radii (mass).

To ascertain the validity of this claim, we first briefly discuss the relevance of detection biases in the list of KOIs. First, one might expect a deficiency of transits at high impact parameters (i.e., $\langle b \rangle$ less than 0.5) due to the lower S/N for shorter transit durations, and due to the removal of V-shaped grazing transits from the list of KOIs because of increased probability of false-positives from stellar eclipsing binaries. B12 includes estimates of the impact parameter b , which can be determined from the ingress and egress times for high S/N transits after accounting for limb-darkening (Seager & Mallen-Ornelas 2003). This detection bias is only readily apparent for KOIs with $b > 0.9$ as tabulated in B12, and it is not correlated with exoplanet radius. In fact, because a number of the KOIs are eccentric and transiting near periastron (as opposed to circular orbits), the distribution of calculated impact parameter in B12 has a median value of 0.7, which is greater than 0.5. This larger value for $\langle b \rangle$ suggests both that Kepler can adequately detect short duration transits, and that some of the KOIs must be on eccentric orbits, further exasperating the discrepancy between the observed mean value of α for all KOIs and that expected for a population of exoplanets on circular orbits with uniform random impact parameters. We do not include the

effects of limb-darkening on our analysis. We also do not use the impact parameters in B12 since a circular orbit is assumed, and there are large uncertainties in constraining b due to the Kepler 30-minute cadence (Fabrycky 2012). Regardless, we find that adjusting the upper bound to the assumed impact parameter distribution in our analysis does not alter our results.

Additionally, one might expect that there is a deficiency of small exoplanets with short transit durations relative to larger exoplanets and/or longer transit durations at higher S/N, leading to a bias towards larger α values for smaller exoplanets. However, this bias does not appear to be present for the KOI parameters tabulated in B12. All but 7 KOIs have a transit detection S/N > 7, and there is no observed trend of transit duration in hours as a function of planet radius. This lack of a bias in our analysis is due to the long Kepler baseline, and the fact that transits are longer for longer orbital periods. However, we do not draw any conclusions regarding the frequency or distribution of exoplanets as a function of orbital period, which would necessarily have to take this bias into account.

4. Comparison to Radial Velocity Discovered Exoplanet Eccentricities

Since the impact parameters and angles of periastron for KOIs are generally not constrained, Equation 2 does not permit a direct quantitative statistical comparison to the eccentricity distribution of RV discovered exoplanets. As recognized by Burke (2008), F08 and others, the distribution of transit duration anomalies α can be used as a proxy instead. We convert the distribution of eccentricities of confirmed RV exoplanets into a simulated distribution of α transit duration anomalies. We assume that every radial velocity exoplanet “transits” with a distribution of impact parameters b and angles of periastron ω that are properly weighted by the transit probability. As noted previously, the angle of periastron cannot be assumed to be uniform random on a sphere as is true for RV discovered exoplanets. An accurate probability distribution for the impact parameter, eccentricity, and angle of periastron for transiting planets is given in Burke (2008, Equations 14–16; Figure 4). We numerically integrate the transit probability using the prescription in (Burke 2008) combined with the known RV sample eccentricity values to derive a simulated distribution of α values for the RV sample. We then directly compare this to the CDFs of α values for all KOIs, KOIs with $R_{pl} < 2R_{\oplus}$, $R_{\oplus} < R_{pl} < R_{\oplus}$, and $R_{pl} > 6R_{\oplus}$. We plot these CDFs in Figure 1.

The simulated RV sample α distribution can be approximated by a normal distribution with a mean and standard deviation of $(\mu, \sigma) = (0.714, 0.305)$ ($\chi^2=0.659$). This result is consistent with the models in F08 for an assumed Rayleigh distribution of eccentricities with an average eccentricity of ~ 0.2 – 0.3 . This also verifies that we correctly computed the α distribution from the eccentricities. A large value of $\alpha > 1$ (assuming accurate stellar parameters) requires a high eccentricity and transit near apastron, which is a low probability occurrence. This explains the notable lack of simulated transit duration anomalies with $\alpha > 1.5$, which is consistent with Burke (2008, Figure 4).

The α distribution for all KOIs with $(\mu, \sigma) = (0.921, 0.288)$ ($\chi^2=2.03$) is significantly different from the RV α distribution, in disagreement with Kane et al. (2012). We calculate distribution mean and standard deviations of $(\mu, \sigma) = (0.978, 0.255)$ for $R_{pl} < 2R_{\oplus}$, $(0.901, 0.279)$ for $2R_{\oplus} < R_{pl} < 6R_{\oplus}$, and $(0.761, 0.331)$ for $R_{pl} > 6R_{\oplus}$ ($\chi^2=0.669, 1.02, 0.556$). The differences between almost all of these distributions are statistically significant. Formally, we evaluate the 1-D Kolmogorov-Smirnov (K-S) test statistic between all distributions. The α distribution for RVs has a probability of being drawn from the same parent population as the KOIs with $R_{pl} > 6R_{\oplus}$ ($2R_{\oplus} < R_{pl} < 6R_{\oplus}, R_{pl} < 2R_{\oplus}$, all) of 4.2×10^{-4} ($4.0 \times 10^{-23}, 4.7 \times 10^{-40}, 1.5 \times 10^{-30}$). The KOIs with $2R_{\oplus} < R_{pl} < 6R_{\oplus}$ and $R_{pl} < 2R_{\oplus}$ have a probability of 13% of being drawn from the same parent population ($< 2\sigma$) and are thus the most similar, whereas the KOIs with $R_{pl} > 6R_{\oplus}$ are statistically distinct from the KOIs with $2R_{\oplus} < R_{pl} < 6R_{\oplus}$ and $R_{pl} < 2R_{\oplus}$ with K-S test probabilities of 3.0×10^{-9} and 1.4×10^{-19} respectively ($> 5\sigma$). In other words, KOIs with $R_{pl} > 6R_{\oplus}$ most closely resemble the RV distribution perhaps as expected, but are still statistically distinct at the $> 3\sigma$ level (and with higher statistical significance for the remaining KOIs). This is also visually apparent from Figure 1. Similarly, the KOIs with $R_{pl} < 6R_{\oplus}$ appear to be drawn from a distinct eccentricity distribution than KOIs with $R_{pl} > 6R_{\oplus}$, with little difference, if any, above and below $R_{pl} = 2R_{\oplus}$.

From this we can conclude that smaller radius KOIs are increasingly more likely to have smaller average eccentricities within these samples. This apparent trend may solely be due to the incompleteness of Kepler at long orbital periods for smaller exoplanets (§3) combined with the well-established tidal circularization of exoplanets at short orbital periods (Butler et al. 1997; Rasio & Ford 1996). Alternatively, Fabrycky et al. (2012) notes that smaller exoplanets are more likely to be found in multiple exoplanet co-aligned systems; the smaller relative mutual inclinations implies smaller average eccentricities.

Finally, the $\alpha < 0.75$ distribution of KOIs with $R_{pl} > 6R_{\oplus}$ most closely (and encouragingly) resembles the RV exoplanet distribution, as might be expected. However, there is a higher occurrence rate of KOIs with $\alpha > 1.5$ than expected for both a Rayleigh eccentricity distribution (F08) and for our simulated RV sample distribution. This appears to be especially true for Jovian KOIs with $R_{pl} > 6R_{\oplus}$. This excess was also noted in Moorhead et al. (2011) and Wang & Ford (2011) and we confirm that this excess is still present with the revised stellar parameters and new list of KOIs. If we assume that our RV and KOI samples are drawn from the same population of exoplanets, some of these KOIs must be attributable to errors in the host stellar parameters or alternatively are false-positives. It is not clear how the stellar radius (mass) could be under-(over-)estimated by factors exceeding 1.5 (1.5^3) to correct some of these α -values. However, a blanket rejection of these KOIs is not recommended since that could inadvertently exclude a genuine highly eccentric exoplanet transiting near apastron.

5. Accuracy of Stellar Parameters Inferred from Transit Duration

The significant difference between the transit duration anomaly α distributions for RV exoplanets and Kepler candidates raises the question about the validity of the KOI host stellar parameters. In Figure 2 we plot the dimensionless α parameter as a function of stellar mass, radius and effective temperature, for both the second and current lists of KOIs. We bin the α values for each stellar parameter, with bins of 250 K in effective temperature, $0.1 M_{\odot}$ in stellar mass, and $0.1 R_{\odot}$ in stellar radius. Our results are insensitive to adjustments in the bin width. We derive median values and quartile ranges for each bin, which are over-plotted in Figure 2 and show distinct trends as a function of stellar mass and radius for both the old and new KOI lists. To quantify these trends, we fit a line to the median values, excluding bins with fewer than 4 KOIs.

The new list of KOIs yields less scatter in the median values of α as a function of stellar parameters, indicating that the revised stellar parameters in B12 are indeed improved. However, we identify a statistically significant trend for decreasing stellar radius (mass) of a 0.27 (0.28) change in α per $R_{\odot}$ ($M_{\odot}$) at the $\sim 6\text{-}\sigma$ ($3\text{-}\sigma$) level for both the current and old KOI lists. The ensemble α values are not found to be linearly biased as a function of stellar effective temperature. The identified trends in mass and radius may have been overlooked in Moorhead et al. (2011) because they only examined the CDFs of the α distribution as a function of T_{eff} . Additionally, the linear trend in stellar temperature for the old KOI list is statistically significant if one excludes KOIs with $T_{eff} > 7000$ K where the median α values decrease again (not shown in Figure 2). This trend is not seen for the new KOI stellar effective temperatures.

The trends in stellar mass and radius imply one of two scenarios. First, there could be a systematic error in these two parameters for the ensemble of KOIs. If we assume that the median α value for $1.0 R_{\odot}$ ($M_{\odot}$) KOIs is accurate, then the ensemble radii (or masses) are under- (over-)estimated by $\sim 15\%$ ($\sim 50\%$) at $0.5 R_{\odot}$ ($M_{\odot}$), and the ensemble radii (or masses) are over- (under-)estimated by $\sim 15\%$ ($\sim 50\%$) at $1.5 R_{\odot}$ ($M_{\odot}$). The larger implied error in stellar mass is due to the weaker $1/3$ -rd power dependence of α on stellar mass. The ensemble stellar masses in B12 and the KIC are likely not in error by $\sim 50\%$ at the low- and high- mass ends. The α -trend bias is thus largely due to systematic ensemble errors in the stellar radius at the low- and high-radius ends by the indicated amounts.

The first scenario relies on the assumption that the eccentricity distribution of exoplanets is intrinsically stellar-mass independent. Alternatively, Howard et al. (2011) suggests that terrestrial planets are more common around lower-mass stars, and Figure 1 demonstrates that terrestrial planets may preferentially be found on more circular orbits. This picture would at first appear to be consistent with the observed trends in α both as a function of stellar mass and radius. In other words, the systematic “error” in α could instead be due to a real change in eccentricity distributions. A $\sim 15\%$ change in α between 0.5 and $1.0 M_{\odot}$ (or 1.0 and $1.5 M_{\odot}$) would correspond to a > 0.2 change in the average ensemble eccentricity of $e \sim (0, 0.25, 0.6)$ for $(0.5, 1, 1.5) M_{\odot}$ as inferred from Figure 9 in F08. Such a large trend in average eccentricity is not observed for the RV exoplanets

in our sample.

Unfortunately, these two scenarios are degenerate, so we cannot rule out a modest change in ensemble eccentricity distributions as a function of stellar mass. However, given the lack of a large eccentricity dependence on stellar mass for RV discovered exoplanets, we conclude that the bulk trends in ensemble α as a function of stellar mass and radius for KOIs are likely entirely due to systematic errors in stellar radii. The ensemble KOI under-estimate of stellar radii for K/M dwarf runs contradictory to the results of Muirhead et al. (2012).

Finally, the over-abundance of α values greater than 1.5 identified in §4 is independent of stellar mass, radius and temperature. We compare KOI α values against every property calculated in the KOI tables in B12. We find that KOIs with $\alpha > 1.5$ are preferentially found around low metallicity KOIs with $\log[Fe]/[H] < -0.11$ for all planet radii, and for $A_V > 0.33$ mag for $R_{pl} > 6R_{\oplus}$ as shown in Figure 3. The cutoff values for $\log[Fe]/[H]$ and A_V correspond to the sample medians. While the extinction may be a crude proxy for brightness, there is no a priori reason to expect these two trends. Thus we conclude that the over-abundance of KOIs with $\alpha > 1.5$ is due to errors in the calculated stellar metallicities (and possibly extinction) for these sources, rather than constituting a genuine class of exoplanets with high eccentricities (improbably) transiting far outside of periastron. These errors in turn imply under-estimates of the stellar radii for these KOIs of $>50\%$. Instead, these stellar hosts are likely more distant (sub-)giants with larger stellar radii and larger secondary companions.

6. Conclusions

We have carried out an updated analysis of the transit duration anomalies with the recently released list of new Kepler exoplanet candidates. In particular, we looked at the KOI distribution of transit durations compared to what would be expected from the eccentricity distribution of RV discovered exoplanets as a function of stellar host parameters. We find systematic errors in the stellar radius and metallicity preclude a quantitative ensemble comparison of the two samples at this time. However, we find that terrestrial planets may preferentially be found on more circular orbits compared to Jovian planets. The systematic bias in stellar radius impacts the inferred distributions of exoplanet properties, including radius and habitability.

The authors would like to thank the anonymous referee for their comments and suggestions. The authors thank John Johnson, Dan Fabrycky, Philip Muirhead, Andrew Youdin, and Alex Hayes for their useful discussions. This research has made use of the NASA Exoplanet Archive, which is operated by the California Institute of Technology, under contract with the National Aeronautics and Space Administration under the Exoplanet Exploration Program.

REFERENCES

- Barnes, J.W., 2007, PASP, 119, 986
- Batalha, N., et al., 2012, ApJS, submitted, arXiv 1202.5852
- Batalha, N., et al., 2010a, ApJL, 713, 109
- Batalha, N., et al., 2010b, ApJL, 713, 103
- Borucki, W., et al., 2012, ApJ, 745, 120
- Borucki, W., et al., 2011a, ApJ, 728, 117
- Borucki, W., et al., 2011b, ApJ, 736, 19
- Borucki, W., et al., 2010, Science, 327, 977
- Brown, T., et al., 2011, AJ, 142, 112
- Burke, C., 2008, ApJ, 679, 1566
- Butler, P., et al., 1997, ApJL, 474, 115
- Ciardi, D., et al., 2011, AJ, 141, 108
- Dawson, R., & Johnson, J., 2012, ApJ, submitted, arXiv 1206.5579
- Fabrycky, D., 2012, priv. comm.
- Fabrycky, D., et al. 2012, ApJ, submitted, arXiv 1202.6328
- Figueira, P., et al., 2012, A&A, submitted, arXiv 1202.2801
- Ford, E., Quinn, S., & Veras, D., 2008, ApJ, 678, 1407
- Howard, A., et al., 2011, ApJ, submitted, arXiv 1103.2541
- Kane, S., Ciardi, D., Gelino, D., & von Braun, K., 2012, MNRAS, in press, arXiv 1203.1631
- Mann, A., 2012, ApJ, submitted, arXiv 1202.5394
- Morton, T., & Johnson, J., 2011, ApJ, 729, 138
- Moorhead, A., et al., 2011, ApJS, 197, 1
- Muirhead, P., et al., 2012, ApJ, 747, 144
- Muirhead, P., et al., 2011, ApJL, submitted, arXiv 1109.1819
- Pinsonneault, M., et al., 2012, ApJS, in press, arXiv 1110.4456

- Plavchan, P., & Bilinski, C., 2012, ApJ, submitted, arXiv 1112.1595
- Rasio, F. A., & Ford, E. B., 1996, Science, 274, 954
- Seager, S., & Mallen-Ornelas, G., 2003, ApJ, 585, 1038
- Steffen, J., et al., 2010, ApJ, 725, 1226
- Traub, W., 2012, ApJ, 745, 20
- Youdin, A., 2011, ApJ, 742, 38
- Wang, Ji., & Ford, E., 2011, MNRAS, 418, 1822

Table 1. KOI Transit Duration Anomaly and Eccentricity Limits

KOI	α	e_p^a	e_a^b
1.01	0.589	0.485	...
2.01	0.666	0.386	...
3.01	0.903	0.102	...
4.01	0.444	0.671	...
5.01	0.467	0.642	...
5.02	0.776	0.248	...
7.01	1.22	...	0.198
10.01	0.710	0.330	...
12.01	1.12	...	0.117
13.01	0.658	0.396	...

Full table available online in electronic format.

^aFor a transit assumed to occur at periastron with a zero impact parameter.

^bFor a transit assumed to occur at apastron with a zero impact parameter.

Table 2. Candidate Eccentric KOIs

KOI	α	e_p^a	e_a^b
1845.02	0.155	0.953	...
371.01	0.156	0.952	...
2519.01	0.178	0.939	...
2522.01	0.190	0.931	...
2287.01	0.199	0.924	...
403.01	0.206	0.918	...
1815.01	0.221	0.907	...
1164.01	3.90	...	0.877
2046.01	2.47	...	0.719
783.01	2.41	...	0.7077

^aFor a transit assumed to occur at periastron with a zero impact parameter.

^bFor a transit assumed to occur at apastron with a zero impact parameter.




Fig. 1.— Cumulative Distribution Functions for: the distribution of transit duration anomalies α modeled from the distribution of RV-discovered exoplanet eccentricities (black) with a mean and standard deviation of $(\mu, \sigma) = (0.714, 0.305)$, the distribution of α for all KOIs (red) with $(\mu, \sigma) = (0.921, 0.288)$, $R_{pl} < 2R_{\oplus}$ KOIs (blue) with $(\mu, \sigma) = (0.978, 0.255)$, $2R_{\oplus} < R_{pl} < 6R_{\oplus}$ KOIs (green) with $(\mu, \sigma) = (0.901, 0.279)$, and $R_{pl} > 6R_{\oplus}$ KOIs (pink) with $(\mu, \sigma) = (0.761, 0.331)$.




Fig. 2.— Distributions for the transit duration anomalies α as a function from left to right of stellar radius, mass, and effective temperature for the new KOI list in the top row, and old KOI list in the bottom row, respectively. Overplotted are binned median, lower and upper quartile ranges, with bin sizes of $0.1 R_{\odot}$, $0.1 M_{\odot}$, and 250 K. Some bins are missing upper/lower quartile ranges since only a couple stars fall into that particular bin. Linear fits to the binned median values (not shown) yield slope coefficients clockwise from top left of -0.27 ± 0.05 , -0.28 ± 0.09 , $-(3.2 \pm 2.6) \times 10^{-5}$, $-(8.3 \pm 4.7) \times 10^{-5}$, -0.28 ± 0.09 , & -0.33 ± 0.05 respectively with the appropriate units, indicating statistically significant ensemble trends for stellar radius and mass.




Fig. 3.— Distributions for the transit duration anomalies α as a function of stellar metallicity on the left and extinction on the right. KOIs with exoplanet radii of $R_{pl} < 2R_{\oplus}$, $2R_{\oplus} < R_{pl} < 6R_{\oplus}$, and $R_{pl} > 6R_{\oplus}$ are shown in blue, green and red respectively. The horizontal line at $\alpha = 1.5$ corresponds to a CDF of $\sim 100\%$ for simulated α values from the known eccentricity distribution of RV-discovered exoplanets. This indicates that most $\alpha > 1.5$ KOIs likely have large systematic errors in stellar parameters. The vertical lines correspond to the median values for all KOIs in metallicity (-0.11) and extinction (0.33).